



Contribution ID: 77

Type: **Poster presentation**

## **The impact of sudden stratospheric warmings (SSWs) on stratosphere-troposphere exchange of ozone (O<sub>3</sub>) and water vapour (H<sub>2</sub>O)**

Midwinter sudden stratospheric warmings (SSWs), characterised by the reversal of the temperature gradient poleward of 60°N and the 10 hPa climatological zonal mean wind from westerly to easterly at 60°N, are known to have pronounced impacts on tropospheric circulation which lead to regional changes in temperature, precipitation and other meteorological variables. Such abrupt events are furthermore known to be associated with large-scale changes in the distribution of stratospheric chemistry constituents, such as ozone (O<sub>3</sub>) and water vapour (H<sub>2</sub>O), although the implications for stratosphere-troposphere exchange (STE) have not been previously investigated. The evolution of O<sub>3</sub> and H<sub>2</sub>O anomalies during an SSW life cycle are first examined from the surface up to 1 hPa using specified-dynamics simulations from the European Centre for Medium-Range Weather Forecasts –Hamburg (ECHAM)/Modular Earth Submodel System (MESSy) Atmospheric Chemistry (EMAC) model over the period 1979-2013. We show that significant positive anomalies in O<sub>3</sub> occur around the onset of an SSW in the middle to lower stratosphere, with persistence timescales of around 50 days in the upper troposphere-lower stratosphere (UTLS). Similarly, we find significant H<sub>2</sub>O anomalies in the lowermost stratosphere ( $\pm 25\%$ ) for up to 75 days. The extent and magnitude of the anomalies are largely confirmed in both Copernicus Atmospheric Monitoring Service (CAMS) reanalysis and ozonesonde measurements at five different Arctic stations. Evaluation of the vertical residual velocity ( $w^*$ ) support the notion of transport changes being the driver of the temporal evolution of the anomalies. Using a stratospheric-tagged O<sub>3</sub> tracer, a signal for enhanced STE of ozone is subsequently inferred ( $\sim 5\text{-}10\%$ ), which is maximised around 50 days after the SSW onset date. Our results highlight that SSWs can induce significantly disturbed O<sub>3</sub> and H<sub>2</sub>O distributions in the UTLS, leading to enhanced STE of O<sub>3</sub>, with potentially significant implications for radiative forcing and air quality.

Key words: Sudden stratospheric warmings (SSWs), upper-troposphere lower-stratosphere (UTLS), stratosphere-troposphere exchange (STE), ozone (O<sub>3</sub>), water vapour (H<sub>2</sub>O)

**Primary author:** Mr WILLIAMS, Ryan (University of Reading)

**Co-authors:** HEGGLIN, Michaela I (University of Reading); JÖCKEL, Patrick (German Aerospace Center (DLR), Institute of Atmospheric Physics, Oberpfaffenhofen-Wessling, Germany); GARNY, Hella (German Aerospace Center (DLR), Institute of Atmospheric Physics, Oberpfaffenhofen-Wessling, Germany)

**Presenter:** Mr WILLIAMS, Ryan (University of Reading)

**Session Classification:** Poster session

**Track Classification:** Workshop: Stratospheric predictability and impact on the troposphere