


New Coupled Modeling System based on the Korean Integrated Model (KIM)

Myung-Seo Koo, Eunjeong Lee, Mee-Hyun Cho, Hyeon-Ju Gim, Junseong Park, Jaeyoung Song, Yong-Jae Han, Jin-Yun Jeong, Subin Kim, and Duho Jeong Korea Institute of Atmospheric Prediction Systems

1 Background

- The Korean Integrated Model (KIM) was developed for global numerical weather prediction and has been operational in real-time since April 2020.
- To expand its capabilities from deterministic medium-range weather forecast to probabilistic sub-seasonal to seasonal prediction, the Korea Institute of Atmospheric Prediction Systems (KIAPS) is developing a new coupled modeling system by advancing the land surface processes and coupling the ocean, sea ice, wave, and river-routing models, along with the further development in ensemble forecast and coupled data assimilation.
- This presentation will provide an overview, evaluation result, and future plan of the KIM-based coupled modeling system.


2 Framework of coupled KIM




3 Configuration of surface models

KIM-CPL		LSM			OCN	SIC	WAV	RIV
Model		Noah-MP			NEMO	SI ³	WaveWatch III	CaMa- Flood
	current	4.0.1 4.2 → 4.4		5.0	4.0		7.13	4.0.0 (4.1)
Ve	c. origin	LIS	WRF	GitHub		-	-	-
	latest	5.0			5.0 beta		7.14	4.2
	Coupler	-			MCT-based			
Initial data		ERA5			ORAS5 (GODAPS)	ERA5/GIOMAS PIOMAS(N.H.)	-	-
Exchange frequency Grid system		every time step (same with KIM)			1h (fixed; same with radiation)			
		cubed-sphere (same with KIM)			tripolar		(regular) lat-lon	unit- catchment
Resolution		100~6 km (NE045~768L91; same with KIM)			25 km (eORCA025L70; fixed)		50 km (fixed)	25 km (fixed)

4 Results





5 Summary and future plan

KIM coupled modeling system

- has been newly developed by coupling the state-of-the art surface model.
- is being updated and provides deterministic forecast skill comparable to that of the operational KIM, along with better performance on long-term simulations.
- shows potential to help improve medium- and long-term forecasting performance. In the future, we will
- adopt more sophisticated physics/model (lake, ...) and realistic data (LAI, ...).
- employ (weakly) coupled data assimilation system by utilizing LIS and NEMO-VAR.
- deeply evaluate on sub-seasonal (MJO, ...) to decadal (ENSO, ...) prediction.