

# Using the Atmosphere-Ocean Single-Column Model (AOSCM) with HALO-AC3 and other observational campaigns to understand model representation of warm and moist air intrusions to the Arctic

Gunilla Svensson,  
Michail Karalis and Michael Tjernström




# Why Lagrangian and Eulerian frameworks?

- Eulerian frameworks are useful to evaluate models at locations
- However, air mass transformation happens along the transport pathway
- Many studies of their importance for the climate are based on models or reanalysis (e.g. Woods et al. 2013, Murto et al. 2022; 2023)
- We know that models and reanalysis have problems representing processes such as turbulent mixing and cloud properties (e.g. Tjernström et al., 2021)



Warm air intrusions/  
atmospheric rivers

Cold air outbreaks




# Why Lagrangian and Eulerian frameworks?

- Physical processes of importance are radiative cooling, cloud formation and turbulent mixing, surface exchange
- Aim here is to build a modeling framework that can be used to improve representation of the parameterized small-scale processes



# Can we utilize the single-column modeling framework?




Same model components as IFS/OpenIFS, EC Earth and ERA5/ERA6



*Atmosphere-ocean single-column model (AOSCM)*  
*Hartung et al., 2018, 2022*


# Can we utilize the single-column modeling framework?

Decoupling the large-scale circulation and the effect of the parameterized small-scale processes makes it possible to investigate process representation without changing the large-scale (resolved) flow




**Eulerian view**

# Can we utilize the single-column modeling framework?



Lagrangian view

De-coupling the large-scale circulation and the effect of the parameterized small-scale processes makes it possible to investigate process representation without changing the large-scale (resolved) flow



Eulerian view


OpenIFS  
SCM cy43r3

OASIS cpl


NEMO3.6/LIM  
or NEMO4/ SI<sup>3</sup>

# HALO-AC3 campaign provides observations to test the framework



12 March 2022, 12UTC




13 March 2022, 12UTC





14 March 2022, 12UTC




# Experimental setup for the Halo-AC3 warm air intrusion





# Experimental setup for the Halo-AC3 warm air intrusion



- Advective tendencies = **off**
- **Varying SST** over ocean,
- Coupled to sea-ice model over sea-ice
- Prescribed ERA5 vertical velocity  $\omega$  and **nudged to ERA5 wind** throughout the column




# HALO-AC3 campaign provide observations to test the framework




# AOSCM simulated tendencies for temperature and moisture



— ocean    - - - MIZ    — ice



The tendency analysis shows when and where processes are important for the transformation






# Expanding to other warm air intrusions



MOSAiC, 19/02  
HALO-AC3, 13/03  
HALO-AC3, 15/03  
MOSAiC, 16/04  
AoM, 12/06  
ACSE, 07/08  
MOSAiC, 13/09  
MOSAiC, 22/09  
MOSAiC, 16/11  
MOSAiC, 03/12


48 h trajectories  
Ensemble simulations  
for 6 pressure levels  
spanning 500 – 900 hPa

# Expanding to other warm air intrusions



# Expanding to other warm air intrusions


- Similar evolution in AOSCM and ERA5
- Air masses reach saturation over deep (5 km) layers
- Vertical ascent is a key driver for the transformation
- How sensitive is the transformation to parameterized processes?



# Perturbations are evaluated on their effect on the energy balance at the top of the atmosphere (TOA) and the surface (SEB)

Top of the atmosphere


$\tilde{\text{SEB}}, \times 1$   
 $\tilde{\text{TOA}}$   
 $\times 1/10$   
 $\times 1/100$



Surface

# Large seasonal differences in the TOA and SEB response of microphysical parameters such as the cloud droplet number concentration


Top of the atmosphere



Surface


# Perturbations in the boundary layer mixing properties drive considerable changes in the SEB in all seasons

Top of the atmosphere




Surface


# Framework summary




A Lagrangian single-column modeling framework can realistically simulate the Arctic air mass transformation



Large-scale ascent is the dominant driver of the transformation in ten WAIs across the seasonal cycle



The tendency analysis can be analyzed per timestep and process



The energy impact of Arctic air mass transformation is sensitive to perturbations in cloud and boundary layer processes

## Some final thoughts...

- AOSCM simulations are similar enough to the full model to be useful for examining the processes and their representation
- Interaction with the sea ice important for the evolution
- Vertical velocity is very important for the transformation, at least in the model
- HALO-AC3 observations revisited the same airmass, but only a limited part of the transformation was covered
- Framework has the potential to be used to design observational campaigns – upcoming CONIDA + IB Oden?
- Can be used, in combination with the full global model, to test and develop parameterizations

Thanks for listening! Questions?

## Arctic air mass transformation

A Lagrangian view of the dynamical and physical drivers

Michail Karalis

