



Coupled land-atmosphere variability and predictability

Jonny Day / Bart van den Hurk / Tim Stockdale


jonathan.day@ecmwf.int



Is the atmosphere heated from above or below?



Earth's energy budget



Why do we care about land processes?


- Energy-budget
 - Albedo




Surface	Albedo
Dark forest	9-12%
Grassland	15-20%
Bare soil	20-30%
Snow in forest	15-25%
Open snow	50-85%

Example of snow transitions

Snow and albedo




Snow and surface meteorology



Betts et al., (2014)

Why do we care about land processes?


- Energy-budget
 - Albedo
 - Evaporative fraction



Surface	LE/Q^*
Boreal forest	25%
Forest in temperate climate	65%
Dry vineyard	20%
Irrigated field in dry area	100%


Why do we care about land processes?

- Energy-budget
 - Albedo
 - Evaporative fraction
- Water budget
 - Runoff-fraction




Land processes in atmospheric models

- Energy-budget
 - Albedo
 - Evaporative fraction
- Water budget
 - Runoff-fraction
 - Soil water reservoir




Land processes in atmospheric models

- Energy-budget
 - Albedo
 - Evaporative fraction
- Water budget
 - Runoff-fraction
 - Soil water reservoir
- Momentum budget
 - Roughness elements




Land processes in atmospheric models

- Energy-budget
 - Albedo
 - Evaporative fraction
- Water budget
 - Runoff-fraction
 - Soil water reservoir
- Momentum budget
 - Roughness elements
- Carbon budget
 - Not directly relevant for seasonal forecasting, but vegetation changes have feedbacks on other processes and are important for climate modelling




What is needed to contribute to predictability?

- In the climate system all processes are connected



- A systematic influence of land surface on atmosphere requires:
 - **Variability**
 - **Memory**
 - **Coupling to the atmosphere**



Otherwise: Can just use downstream/application models, e.g. crop modelling, hydrology flood forecasting, fire risk models etc

Soil moisture predictability

$$c_s w_{n+lag,y} = c_s w_{n,y} + P_{n,y} - E_{n,y} - Q_{n,y}$$

$$\rho(w_n, w_{n+lag}) = \frac{cov(w_n, w_{n+lag})}{\sigma_{w_n} \sigma_{w_{n+lag}}}$$


w_n =soil moisture at time n

P_n =total precipitation between time n and $n+lag$


E_n =total evaporation between time n and $n+lag$

Q_n =total runoff between time n and $n+lag$


C_s =water holding capacity



Spatial variation in persistence and skill of root-zone soil-moisture



Seasonal forecast anomaly correlation JJA




Day et al., 2025



Funded by the
European Union

Soil-moisture atmosphere coupling



$$E = \beta E_{POT} = \beta \rho_a \left[\frac{q_{sat}(T_s) - q_r}{r_a} \right]$$

with

$$\beta = \frac{\theta - \theta_{WILT}}{\theta_{CRIT} - \theta_{WILT}} \text{ for } \theta_{WILT} \leq \theta \leq \theta_{CRIT}$$

$$\beta = 1 \text{ for } \theta > \theta_{CRIT}$$


$$\beta = 0 \text{ for } \theta < \theta_{WILT}$$

Seneviratne et al., 2010

Measures to quantify land-atmosphere coupling


- From (pseudo)observations:
 - Correlation between evaporation and temperature

Seneviratne et al, 2010



Feb-Apr

Corr monthly CTL evap-t2m m02 EOC



Over mid-latitude oceans, evaporation depends on humidity: colder air is generally drier

Over moisture-limited land, drier conditions reduce evaporation and cause higher temperatures

Energy limited


If soils are wet, higher temperatures drive higher evaporation

Soil water limited


Measures to quantify land-atmosphere coupling

- From (pseudo)observations:
 - Correlation between evaporation and temperature

May-Jul




Energy limited



Soil water limited


Seneviratne et al, 2010




Measures to quantify land-atmosphere coupling

- From (pseudo)observations:
 - Correlation between evaporation and temperature

Aug-Oct




Energy limited



Soil water limited


Seneviratne et al, 2010




Measures to quantify land-atmosphere coupling

- From (pseudo)observations:
 - Correlation between evaporation and temperature

Nov-Jan




Energy limited




Soil water limited

Seneviratne et al, 2010




Measures to quantify land-atmosphere coupling

- From a model experiment (GLACE = Global Land Atmosphere Coupling Experiment)
- How?
 - Simulate the hydrological cycle **with** (W) and **without** (R) interactive land-atmosphere coupling and compare.
- How to remove coupling?
 - In second ensemble (R), replace soil moisture in all ensemble members by values from one of the integrations in the first (interactive) ensemble.
- How to measure the effect?
 - Ensemble simulations
 - Compare within-ensemble spread




PART 1: ESTABLISH A TIME SERIES OF SURFACE CONDITIONS (Simulation W1).



(Repeat without writing to obtain simulations W2-W16.)

PART 2: RUN 16-MEMBER ENSEMBLE, WITH EACH MEMBER FORCED TO MAINTAIN THE SAME TIME SERIES OF SURFACE PROGNOSTIC VARIABLES (Simulations R1-R16).




Comparison of precipitation between ensembles

Diagnostics:


$$\Omega = (16\sigma^2_{\langle X \rangle} - \sigma^2_X) / 15\sigma^2_X$$

$\Omega(W)$ = fraction of variance “explained”
(forced) by all boundary and initial conditions

$\Omega(R) - \Omega(W)$ = fraction of variance “explained”
by prescription of subsurface soil moisture
variables



All simulations in ensemble
respond to the specified land surface
boundary condition in the same way
→ strong coupling




Simulations in ensemble
have no coherent response
to the specified land surface
boundary condition
→ weak coupling

Koster et al, 2002

Areas with strong positive soil-moisture feedback on precipitation

Land-atmosphere coupling strength (JJA), averaged across AGCMs




This is a famous figure, and looks very nice. But note that different models gave substantially different results. Model representation of land surface processes is improving, but still has some way to go.


Koster et al, 2004, Science

Compare diagnostic and prognostic measure


Land-atmosphere coupling strength (JJA), averaged across AGCMs





Corr monthly CTL evap-t2m m05 EOC



Strong precipitation coupling needs a combination of sensitivities



Strong feedback on precipitation at transition between arid and temperate zones




Main climates	Precipitation	Temperature
A: equatorial	W: desert	h: hot arid
B: arid	S: steppe	k: cold arid
C: warm temperate	f: fully humid	a: hot summer
D: snow	s: summer dry	b: warm summer
E: polar	w: winter dry	c: cool summer
	m: monsoonal	d: extremely continental
		F: polar frost
		T: polar tundra

Koster et al, 2004, Science


A spatial view on soil-moisture atmosphere coupling

- Chris Taylor et al.:
 - Afternoon rain more likely over drier soils (Nature 2012), apparently at odds with Koster et al. citing errors in convective scale processes in GCMs used by Koster.



Taylor et al. (2011)

Preference for afternoon precipitation over dry soil in obs and a GCM




Taylor et al. (2012)


The Resolution!

- Guillod et al. (2015), Nat. Comms


Koster emphasizes the temporal coupling




Taylor emphasizes the spatial



Remote response to soil-moisture depletion on upper atmosphere




Sensitivity experiment: Impact of North American soil-moisture in JJA 2022 (following a La-Nina)




Difference between phases JJAs following La-Nina vs El-Nino


a) z200 (zonal anomaly) and mean u200





b) soil moisture, ET and SST



How well do GCMs capture soil-moisture-atmosphere coupling?




Snow depth memory




Snow cover as a predictor of the Arctic Oscillation


Corr of DJF AO and DJF SLP 1997-2010




Corr of SAI and DJF SLP 1997-2010



A mechanistic view of remote response to snowcover






From the next model cycle each ECMWF system will have its own land-surface analysis (initial conditions)



Summary

- For land-related predictability we need
 - Variability
 - Memory (soil moisture, snow mass, vegetation, ...)
 - Coupling
- How to measure land-atmosphere coupling/predictability
 - Diagnostic measures: correlation, regression, composite analysis, etc.
 - Prognostic measures: intervening in GCM experiments (GLACE, Teng et al., etc.).
- Land surface signal is small in some regions but large in certain “hotspots”
 - Evaporation limited vs Energy Limited soils
 - Transition zones between semi-arid and humid climates.
- Errors in land-atmosphere feedbacks can lead to large biases that degrade prediction skill on subseasonal-to-centennial timescales.
- Land-surface properties need special treatment (e.g. new LDAS at ECMWF)




References

- Betts, A. K., Desjardins, R., Worth, D., Wang, S., and Li, J. (2014), Coupling of winter climate transitions to snow and clouds over the Prairies, *J. Geophys. Res. Atmos.*, 119, 1118–1139
- Cohen, J., and Jones, J. (2011), A new index for more accurate winter predictions, *Geophys. Res. Lett.*, 38, L21701, doi:10.1029/2011GL049626.
- Day, J.J., Vitart, F., Stockdale, T. et al. Soil-moisture-atmosphere coupling hotspots and their representation in seasonal forecasts of boreal summer. *Clim Dyn* 63, 275 (2025). <https://doi.org/10.1007/s00382-025-07753-1>
- P. A. Dirmeyer, C. A. Schlosser, and K. L. Brubaker, 'Precipitation, Recycling, and Land Memory: An Integrated Analysis', *J. Hydrometeor.*, vol. 10, no. 1, pp. 278–288, Feb. 2009.
- Guillod, B., Orlowsky, B., Miralles, D. et al. Reconciling spatial and temporal soil moisture effects on afternoon rainfall. *Nat Commun* 6, 6443 (2015).
- Henderson, G.R., Peings, Y., Furtado, J.C. et al. Snow–atmosphere coupling in the Northern Hemisphere. *Nature Clim Change* 8, 954–963 (2018).
- B. van den Hurk, F. Doblas-Reyes, G. Balsamo, R. D. Koster, S. I. Seneviratne, and H. Camargo, 'Soil moisture effects on seasonal temperature and precipitation forecast scores in Europe', *Climate Dynamics*, vol. 38, no. 1, pp. 349–362, 2012.
- Koster, R. D., P. A. Dirmeyer, A. N. Hahmann, R. Ijpelaar, L. Tyahla, P. Cox, and M. J. Suarez, 2002: Comparing the Degree of Land–Atmosphere Interaction in Four Atmospheric General Circulation Models. *J. Hydrometeor.*, 3, 363–375
- R. D. Koster et al., 'Regions of Strong Coupling Between Soil Moisture and Precipitation', *Science*, vol. 305, no. 5687, pp. 1138–1140, Aug. 2004.
- R. D. Koster et al., 'Contribution of land surface initialization to subseasonal forecast skill: First results from a multi-model experiment', *Geophys. Res. Lett.*, vol. 37, no. 2, p. L02402, Jan. 2010.
- Koster, R. D., Y. Chang, and S. D. Schubert, 2014: A Mechanism for Land–Atmosphere Feedback Involving Planetary Wave Structures. *J. Climate*, 27, 9290–9301,
- Orth, R., and Seneviratne, S. I. (2012), Analysis of soil moisture memory from observations in Europe, *J. Geophys. Res.*, 117, D15115
- M. Hirschi et al., 'Observational evidence for soil-moisture impact on hot extremes in southeastern Europe', *Nature Geosci*, vol. 4, no. 1, pp. 17–21, Jan. 2011.
- Lin, Y., Dong, W., Zhang, M. et al. Causes of model dry and warm bias over central U.S. and impact on climate projections. *Nat Commun* 8, 881 (2017).
- S. I. Seneviratne et al., 'Investigating soil moisture–climate interactions in a changing climate: A review', *Earth-Science Reviews*, vol. 99, no. 3–4, pp. 125–161, mei 2010.
- Santanello, J. A., and Coauthors, 2018: Land–Atmosphere Interactions: The LoCo Perspective. *Bull. Amer. Meteor. Soc.*, 99, 1253–1272
- CM Taylor RAM Jeu d, Guichard F, Harris P P, Dorigo W A, 2012 Afternoon rain more likely over drier soils *Nature* 489 423 426
- Teng, H., G. Branstator, A. B. Tawfik, and P. Callaghan, 2019: Circumglobal Response to Prescribed Soil Moisture over North America. *J. Climate*, 32, 4525–4545, <https://doi.org/10.1175/JCLI-D-18-0823.1>.


Additional slides

How important is the land-surface?




Trend contributes to T2m predictability

Note: initialized land surface (solid line) gives additional signal in T2m, especially in early summer. Note these plots show predictability not skill – extra skill would require the additional land surface signal to be correct.




Before detrending




After detrending

Impact on T2m forecast skill of improved land surface initial conditions



JJA




DJF

Difference in CRPSS skill from 36 years of re-forecasts with 51 member ensembles, comparing Cy46r1 (ERA5 ICs) and Cy45r1 (SEAS5 ICs). Sensitivity experiments (lower resolution, smaller ensemble size) confirm that the main driver of improvement is change in ICs, not the model.


How about trends in predictability?

- Can we see climate trends in predictability?
 - Model experiment: compare ensemble seasonal forecasts 1900-1929 to 1980-2009
- Can we see trend in land surface contribution to this predictability?
 - Model experiment: same forecasts but with random initial land conditions
- Metric: ratio between **signal** and **total** variance



An anecdote demonstrating impact of soil moisture

- Mid '90's: introduction of **prognostic soil moisture** scheme



- Soil moisture **data assimilation** needed to control drift

(Root cause of drift was model bias, but once unphysical constraint was removed, model bias led to errors that grew over time)

