



Contribution ID: 62

Type: **Oral presentation**

## Flow-dependent predictability of wintertime Euro-Atlantic weather regimes in medium-range forecasts

*Tuesday, 2 April 2019 14:15 (30 minutes)*

This study assesses the medium-range flow-dependent forecast skill of Euro-Atlantic weather regimes: the positive and negative phases of the North Atlantic Oscillation (NAO+ and NAO−), Atlantic ridge (ATLR), and Euro-Atlantic blocking (EABL), for extended winters (November–March) in the periods 2006/2007–2013/2014 and 1985/1986–2013/2014 using The Interactive Grand Global Ensemble (TIGGE) and the National Oceanic and Atmospheric Administration (NOAA)’s Global Ensemble Forecasting System (GEFS) reforecast datasets, respectively. The models show greater-than-observed (smaller-than-observed) frequencies of NAO− and ATLR (NAO+) with forecast lead time. The increased frequency of NAO− is not due to its excess persistence but due to more frequent transitions mainly from ATLR, but also from NAO+. In turn, NAO+ is under-persistent. The models show the highest probabilistic skill for forecasts initialised on NAO− and the NAO− forecasts during the TIGGE period. However, the GEFS reforecast during the period 1985/1986–2013/2014 revealed that these recent high skills reflect the occurrence of four long-lasting (>30 days) NAO− events in 2009/2010–2013/2014 and that the skill for forecasts initialised on NAO− before 2009/2010 (the longest duration was 22 days and the second-longest 16 days) was the lowest. The longer the NAO− events persist, the higher the skill of forecasts initialised on NAO−. The skill dependency on regime duration is less clearly observed for the other regimes. In addition, the GEFS reforecast also revealed that the highest skill of the NAO− forecasts during the period 1985/1986–2013/2014 is attributed to the higher skill of the NAO− forecasts during the active NAO− periods. The EABL forecasts initialised on ATLR show the lowest skill, followed by the NAO− (EABL) forecasts initialised on NAO+ or ATLR (NAO+). These results suggest that the recent models still have difficulties in predicting the onset of blocking.

**Primary authors:** MATSUEDA, Mio (Center for Computational Sciences, University of Tsukuba); Prof. PALMER, Tim (University of Oxford)

**Presenter:** MATSUEDA, Mio (Center for Computational Sciences, University of Tsukuba)

**Session Classification:** Predictability and dynamics

**Track Classification:** Workshop on Predictability, dynamics and applications research using the TIGGE and S2S ensembles